# A novel semi-automated approach for detecting and extracting of urban features using object-based aerial image analysis

- Azra Masumei <sup>1</sup>
- Bakhtiar Feizizadeh <sup>2</sup>
- Khalil Valizadeh Kamran <sup>3</sup>
  - <sup>1</sup> M.Sc. in Remote sensing and GIS, Tabriz University
  - <sup>2</sup> Associate Professor of Remote Sensing and GIS Department of Tabriz University
  - <sup>3</sup> Professor of Remote Sensing and GIS Department of Tabriz University

#### Introduction

Public art has followed a new trend in urban spaces in recent years, so that cities invest highly in the design and implementing of public works of art and improving the visual quality of urban spaces, since it is effective in creating good urban spaces and making the city attractive and memorable. It also encourages people to communicate with works of art and each other. Although public art as an art integrated with the public space of cities is not a new concept, many changes have been created in public art and it caused the creation of various types of public art in most cases from a purely static art to a dynamic and interactive art. This article aims to investigate the effect of public art in urban spaces on promoting the perceived vitality of young people. Despite the importance and role of public art in the quality of urban public spaces, studies conducted in the area of public art have not paid much attention to the effect of different types of public arts in public spaces on vitality of people, especially young people. Thus, this study was conducted with the aim of answering the question of "which type of public art has a greater impact on promoting the perceived vitality of young people in urban spaces?"

## **Data and Methods**

The present study is a qualitative-quantitative (mixed) study conducted as a survey. In the present study, visual approach and image reading were used to collect information. Visual research is a term that includes methods based on data collection and analysis of them through designing, painting, photography, film, or video images. In this article, public arts and interactive arts were divided into different types based on theoretical foundations and research literature and an image was selected from each type of art and through an online questionnaire was submitted to the participants and they were asked to express their level of vitality from watching that work of art in urban spaces on a 5-point Likert scales. This questionnaire was designed and sent to the target population through cyberspace applications. Since the target population of the present article was young people, the sample was selected among the students of Tabriz University and the students of Nabi Akram Institute of Higher Education in Tabriz in the second semester of 2020-2021. To determine the sample size, Cochran's formula with an error level of 5% was used. According to the number of students in these two universities (27000 people), the sample size of 379 was selected. After sending the questionnaire to nearly 380 people, 144 people completed the questionnaire. In this research, both art and architecture students and non-art and architecture students in both bachelor and master levels were surveyed.

## **Discussion and Conclusion**

The research results revealed that among different types of public arts in urban spaces, sound and digital arts such as musical fountain, and the integrated art embedded in the elements available in space such as wall relief in the body of the building and the art appropriate to the site, such as

plants formed in parks, regardless of the three criteria of gender, age, education and field of study, are accepted by all young people and increase vitality, but abstract static art is less welcomed by young people. Among the interactive arts, dynamic-passive interactive art and static interactive art obtained the highest score in terms of creating a sense of vitality among young people, and in contrast, dynamic-dynamic interactive art obtained the lowest score. Art related to public and local culture caused the highest vitality among male young people and young non-art and architecture master students, and digital art caused the highest vitality among females and bachelor art and architecture students. In addition, among non-art and architecture bachelor students, two-dimensional applied art obtained a relatively high score.

#### **Results**

The results show that creating static works of art is not effective in designing cities and using public art to create a sense of vitality in urban space and young people show higher tendency to dynamic work of arts. Also, interactive works of art are more acceptable than non-interactive types, and urban planners should pay attention to this issue in embedding works of art in urban spaces and take right decisions and actions to increase presence, interaction, and vitality among young people in urban spaces. Also, the use of modern technologies and creativity in the artistic design of cities is one of the requirements for adapting the urban space to spirits of young people.

**Key Words:** Urban buildings extraction, Object-based image analysis, accuracy assessment,

Varzaghan city

### **Refrences:**

، بهبود آشکار سازی تغییرات شئ گرا در تصاویر با قدرت تفکیک مکانی بالا بر (1395)اجاقی، سعید ؛ خزائی، صفا ؛ دوره 62، شماره مبنای روش جنگل تصادفی در فضای ویژگی های بهینه ، فصلنامه علمی - بژوهشی اطلاعات جغرافیای ، .401، صص 421-117 ، ار ائه روش تر كيبي از شبكه (1388) حدادي، عطا الله ؛ صاحبي، محمو د رضا؛ مختار زاده ،مهدى ؛ هيرش، فتاحي؛ ایران، دوره 1، GIS های عصبی نظارت شده و نظارت نشده در طبقه بندی تصاویر سنجش از دور، سنجش از دور و شماره 3 ،صص 50-33 عطارچي، سارا ؛ پور رحيمي ، مجتبي ؛ عيسي زاده ، وحيد ؛ (1399)، مقايسهٔ عملكرد شاخص هاي طيفي با طبقه بندی شی ءگرا در استخراج مناطق ساخته شده در مناطق شهری (مطالعهٔ موردی: شهر تهران و شهر گرگان)، پژوهش های . جغرافیای برنامه ریزی شهری، دوره 8، شماره 1، صص43-23 .، مباني سنجش از دور، انتشارات آزاده، تهران(1391) فاطمى، باقر؛ رضايي، بوسف؛ ، مقایسه ی روش های پیکسل پایه و شیءگرا در تهیه نقشه (1386) فیضی زاده، بختیار ؛ رضایی مقدم، محمدحسین ؛ 🛘 دانشگاه تبریز، صص 97-303 GIS های کاربری اراضی، پایانامه کارشناسی ارشد، مرکز فیضی زاده، بختیار ،؛ سلمانی، سعید؛ (1395)، مدلسازی تخریب اراضی کشاورزی بر اثر رشد و توسعه شهری با به 🛘 کارگیری روش های شیءیایه بردازش تصاویر ماهواره ای در محدوده شهری ارومیه،آمایش سرزمین، دوره ۱، شماره 2، .صص 202-177 فیضی زاده، بختیار؛ هلالی، حسین؛ (1388)، مقایسه روش های پیکسل پایه، شیءگرا و پارامتر های تاثیرگذار در طبقه 🛘 بندی پوشش کاربری اراضی استان آذربایجان غربی، پژوهش های جغرافیای طبیعی، شماره 71، صص 84-73. فیضی زاده، بختیار ؛ حاجی میر رحیمی، محمود ؛ (1387)، آشکار سازی تغییر ات کاربری از اضی با استفاده از روش تا 23 ارديبهشت ، سازمان نقشه برداري هاي طبقه بندي شيءگرا مطالعه موردي شهرك انديشه، همايش ژئوماتيك، 22 کشور ، تهران

☐ Aytekin , Ö., Erener, A., Ulusoy, İ. and Duzgun, Ş., 2012.Unsupervised building detection in
complex urban environments from multispectral satellite imagery. International Journal of Remote
Sensing, 33(7): 2152-2177.
☐ Attarzadeh,R.and Momeni,M.:Object-Based Building Extraction from High Resolution
Satellite Imagery, Int. Arch. Photogramm. Remote Sens. Spatial Int. Sci., XXXX-B4,57-60,2012.
□ Blaschke, T. (2010). object based image analysis for remote sensing ISRS Journal of
Photogrammetry and Remote Sensing ,65(1) ,2-16.
☐ Bedawi ,S.and Kamel,M.,2011.Multiple Classifier System for Urban Area Extraction from
High Resolution Remote Sensing Imagery.Image Analysis and Recognition: 307-316.
☐ Congalton,R.G. ,& Green,K.(2008).Assessing the Accuracy of Remotely Sensed Data :
Principles and Practices.2 <sup>nd</sup> ED, Boca Raton,Florida:Lewis Publishers.
☐ Comert, R., Kaplan, O. (2018). Object based building extraction and building period
estimation from unmanned aerial vehicle data. ISPRS Journal of Photogrammetry and Remote
Sensing, 6(3):71-76.
☐ Dervisoglu, Adalet., Bilgilioglu, Baha., Yagmur, Nur.(2020). Comparison of Pixel-Based
and Object-Based Classification Methods in Determination of Wetland Coastline . International
Journal of Environment and Geoinformatics, 7(2): 220-213.
☐ Dragut, L., Eisank, C. (2012). Automated object-based classification of topography from SRTM
data, Geomorphology 141-142,21-33.
☐ Duro, D. C., Franklin, S. E. and Duba, M. G., 2012. A comparison of pixel-based and object-
based image analysis with selected machine learning algorithms for the classification of
agricultural landscape using SPOT-5 HRG imagery.Remote Sensing of Environment,118(0):259-
272.
☐ Erener,A.,2013.Classification method , spectral diversity , band combination and accuracy
assessment evaluation for urban feature detection.International Journal of Applied Earth
Observation and Geoinformation, 21:397-408.
☐ Foody, G. M.,2002, Status of land cover classification accuracy assessment, Remote Sensing of
Enviroment, 80, 185-201.
☐ Feizizadeh , B.,2017. A Novel Approach of Fuzzy Dempster-Shafer Theory for Spatial
Uncertainty Analysis and Accuracy Assessment of Object-Based Image Classification, IEEE
Geoscience and Remote Sensing Letters.
Grinias, I., Panagiotakis, C., Tzirita., G., 2016. MRF-based Segmentation and unsupervised
classification for building and road detection in peri-urban areas of high-resolution Satellite
images. ISPRS Journal of Photogrammetry and Remote Sensing, 122(2016), 145-166.
☐ Huang , D., Shujuei , X., Jingqi , S., Saling , L., Wei , S., Zhenhua , W., 2017. Accuracy
assessment model for classification result of remote sensing image based on spatial
sampling. Journal of applied remote sensing ,11(4) ,1-13.
☐ Jahnson , BA., Ma, L., 2020. Image segmentation and object-based image analysis for
environmental monitoring: Recent areas of interest, Researchers views on the future
priorities.Remoe sensing.2020,12(11):1-9.
☐ Ma, L.; Fu, T.; Blaschke, T.; Li, M.; Tiede, D.; Zhou, Z.; Chen, D.; Ma, X(2017). "Evaluation
of Feature Selection Methods for Object-Based Land Cover Mapping of Unmanned Aerial Vehicle
Imagery Using Random Forest and Support Vector Machine Classifiers", ISPRS International
Journal of GeoInformation, 6(2), 51.

□ Oruc, M., Marangoz, A.M. & Buyuksalih, G. (2004). Comparison of pixel-based and object-
oriented classification approaches using Landsat-7 ETM spectral bands. In Proceedings of XX
ISPRS Congress (p.5), 19 July, Istanbul, Turkey.
□ Otukei , J.R., Blaschke , T., 2010. Land cover change assessment using decision trees, support
vector machines and maximum likelihood classification alghorithms. Supplement Issue on
"Remote Sensing for Africa "A Special Collection from the Africa Association for Remote
Sensing of Environment (AARSE), 12, Supplement 1(0):S27-S31.
□ Pal,M.and Mather,P.M.,2003.An assessment of effectiveness of decision tree methods for land
cover classification.Remote Sensing of Environment,86(4):554-565.
□ Liu,W.and Prinet,V.r.,2005.Building detection from high-resolution Satellite image using
probability model.
☐ Theng,L.B.,2006.Automatic building extraction from Satellite imagery
Engineering, Letters, 13(3):255-259.
☐ Vijjapu, Pavan. (2013), Classification and Building Detection Using Fuzzy Sub_Objects Class
Densities and Shadow Influence, master of science in Geo_information Science and Earth
observation, University of Twente.
•